arXiv:1605.01591v2 [math.GR] 1 Sep 2016

Hall polynomials for the torsion free nilpotent groups of Hirsch length at most 5

Bettina Eick and Ann-Kristin Engel

September 2, 2016

Abstract

A theorem by Hall asserts that the multiplication in torsion free nilpotent groups of finite Hirsch length can be facilitated by polynomials. In this note we exhibit explicit Hall polynomials for the torsion free nilpotent groups of Hirsch length at most 5.

1 Introduction

Let $\mathcal{T}$ denote the class of torsion free finitely generated nilpotent groups. By [7, p. 137], each $\mathcal{T}$-group has a central series with infinite cyclic quotients. The length of such a central series is the Hirsch length of the group; see [7, p. 152] for background.

Let G be a group. We say that a sequence of elements $(g_1, \ldots, g_n)$ is a $\mathcal{T}$-*basis* for G if the subgroups $G_i = \langle g_i, \ldots, g_n \rangle$ for $1 \leq i \leq n+1$ define a central series $G = G_1 > G_2 > \ldots > G_n > G_{n+1} = \{1\}$ with infinite quotients G_i/G_{i+1} for $1 \leq i \leq n$. The construction implies that each quotient G_i/G_{i+1} is cyclic and generated by $g_i G_{i+1}$. It is not difficult to observe that a group G has a $\mathcal{T}$-basis of length n if and only if G is $\mathcal{T}$-group of Hirsch length n.

We note that a $\mathcal{T}$-basis $(g_1, \ldots, g_n)$ of a $\mathcal{T}$-group G is a polycyclic sequence for G in the sense of [5, Sec. 8.1]. Thus as in [5, Lemma 8.3], it follows that each element g of G can be written uniquely as $g = g_1^{a_1} \cdots g_n^{a_n}$ for certain $a_1, \ldots, a_n \in \mathbb{Z}$. Hence the $\mathcal{T}$-basis induces a bijection

$$\beta : \mathbb{Z}^n \to G : (a_1, \ldots, a_n) \to g_1^{a_1} \cdots g_n^{a_n}.$$

The bijection β is not a group homomorphism in general. More precisely, β translates the multiplication in G to certain functions $p_i : \mathbb{Z}^n \times \mathbb{Z}^n \to \mathbb{Z}$ for $1 \leq i \leq n$ with

$$(g_1^{a_1} \cdots g_n^{a_n}) \cdot (g_1^{b_1} \cdots g_n^{b_n}) = g_1^{p_1(a,b)} \cdots g_n^{p_n(a,b)}.$$

A fundamental theorem by Hall [3, 4] asserts that each function $p_i(a, b)$ can be described by a polynomial. These polynomials are nowadays called *Hall polynomials*.

Our aim here is to determine Hall polynomials for *all* $\mathcal{T}$-groups of Hirsch length at most 5. Methods to compute Hall polynomials for a fixed $\mathcal{T}$-group given by a consistent polycyclic presentation have been exhibited by Sims [8, page 441ff] and by Leedham-Green & Soicher [6]. Our method is based on the approach by Sims and uses this in a generic form so that it applies to the infinitely many $\mathcal{T}$-groups of Hirsch length at most 5 simultaneously.

An application of the Hall polynomials determined here is the classification of the torsion free nilpotent groups of Hirsch length at most 5 as described in [2].

2 Nilpotent presentations and consistency

As a first step towards our aim we exhibit consistent nilpotent presentations for the $\mathcal{T}$-groups of Hirsch length at most 5. We refer to [5, Sec. 8.1] for details on this type of presentation. For $n \in \mathbb{N}$ and $t = (t_{ijk} \mid 1 \leq i < j < k \leq n) \in \mathbb{Z}^{\binom{n}{3}}$ we denote

$$G(t) = \langle g_1, \ldots, g_n \mid [g_j, g_i] = g_{j+1}^{t_{i,j,j+1}} \cdots g_n^{t_{i,j,n}} \text{ for } 1 \leq i < j \leq n \rangle.$$

Then the presentation defining $G(t)$ is a polycyclic presentation and the relation imply that $G(t)$ is a nilpotent group of Hirsch length at most n. This presentation is consistent if and only if $G(t)$ has Hirsch length precisely n, or, equivalently, if the generators of the presentation form a $\mathcal{T}$-basis for $G(t)$.

1 Lemma: *Let G be a $\mathcal{T}$-group of Hirsch length 5 and let $(h_1, \ldots, h_5)$ be a $\mathcal{T}$-basis for G. Then there exist a unique $t \in \mathbb{Z}^{10}$ so that the map $G(t) \to G : g_i \mapsto h_i$ extends to an isomorphism. This t satisfies the relations $t_{123}t_{345} = 0$ and $t_{124}t_{345} + t_{145}t_{234} = t_{134}t_{245}$.*

Proof: We first construct a suitable $t \in \mathbb{Z}^{10}$. The definition of $\mathcal{T}$-basis implies directly that $[h_j, h_i] \in G_{j+1}$ for each $1 \leq i < j \leq 5$. As each element of G_{j+1} can be written (uniquely) as $h_{j+1}^{a_{j+1}} \cdots h_5^{a_5}$, this yields that there exist (unique) t_{ijk} with

$$[h_j, h_i] = h_{j+1}^{t_{i,j,j+1}} \cdots h_5^{t_{i,j,5}}$$

for $1 \leq i < j \leq 5$. Using this constructed $t \in \mathbb{Z}^{10}$ then induces that the map $G(t) \to G : g_i \mapsto h_i$ is an epimorphism. It is an isomorphism, as G has Hirsch length 5 and $G(t)$ has at most Hirsch length 5. We now observe that the restrictions on t are satisfied. For this purpose we evaluate the equation $(h_4h_2)h_1 = h_4(h_2h_1)$ in G and obtain that

$$\begin{aligned}
(h_4h_2)h_1 &= (h_2h_4h_5^{t_{245}})h_1 \\
&= h_1h_2^{h_1}h_4^{h_1}h_5^{t_{245}} \\
&= h_1h_2h_3^{t_{123}}h_4^{t_{124}}h_5^{t_{125}}h_4h_5^{t_{145}}h_5^{t_{245}} \\
&= h_1h_2h_3^{t_{123}}h_4^{t_{124}+1}h_5^{t_{125}+t_{145}+t_{245}} \text{ and} \\
h_4(h_2h_1) &= h_4(h_1h_2h_3^{t_{123}}h_4^{t_{124}}h_5^{t_{125}}) \\
&= h_1h_4h_2h_3^{t_{123}}h_4^{t_{124}}h_5^{t_{125}+t_{145}} \\
&= h_1h_2h_4h_3^{t_{123}}h_4^{t_{124}}h_5^{t_{125}+t_{145}+t_{245}} \\
&= h_1h_2h_3^{t_{123}}h_4^{t_{124}+1}h_5^{t_{125}+t_{145}+t_{245}+t_{123}t_{345}}.
\end{aligned}$$

Comparison of the two results yields that $t_{123}t_{345} = 0$. Evaluating the equation $(h_3h_2)h_1 = h_3(h_2h_1)$ in a similar form yields that $t_{124}t_{345} + t_{145}t_{234} = t_{134}t_{245}$. •

As a next step, we show that the restrictions on t in Lemma 1 yield that the presentation defining $G(t)$ is consistent.

2 Lemma: *Let $n \in \{1, \ldots, 5\}$ and $t \in \mathbb{Z}^{\binom{n}{3}}$.*

(a) *Let $n \leq 4$. Then the presentation defining $G(t)$ is consistent.*

(b) *Let $n = 5$. Then the presentation defining $G(t)$ is consistent if and only if $t_{123}t_{345} = 0$ and $t_{124}t_{345} + t_{145}t_{234} = t_{134}t_{245}$.*

Proof: We prove (b) only and note that (a) follows by a similar (and easier) calculation.
$\Rightarrow$: Suppose that $G(t)$ is consistent. Then the generators of this presentation form a $\mathcal{T}$-basis for $G(t)$. Thus Lemma 1 yields the result.
$\Leftarrow$: Suppose that $t \in \mathbb{Z}^{10}$ with $t_{123}t_{345} = 0$ and $t_{124}t_{345} + t_{145}t_{234} = t_{134}t_{245}$ is given. We show that $G(t)$ is consistent in this case. For this purpose we evaluate the finitely many consistency relations, see [8, page 424] (or [1, Lemma 2.10]). In the case considered here, these consistency relations have the form

$$(g_k g_j)g_i = g_k(g_j g_i) \text{ for } 1 \leq i < j < k \leq 5, \text{ and}$$

$$g_k = (g_k g_i^{-1})g_i \text{ for } 1 \leq i < k \leq 5.$$

It is not difficult to observe that these relations always hold if $k = 5$, since g_5 is central in G. Hence it remains to consider the relations

$$\begin{aligned}
(R1) \quad & (g_4g_3)g_1 = g_4(g_3g_1),\\
(R2) \quad & (g_4g_2)g_1 = g_4(g_2g_1),\\
(R3) \quad & (g_3g_2)g_1 = g_3(g_2g_1),\\
(R4) \quad & (g_4g_3)g_2 = g_4(g_3g_2),\\
(R5) \quad & g_2 = (g_2g_1^{-1})g_1,\\
(R6) \quad & g_3 = (g_3g_1^{-1})g_1,\\
(R7) \quad & g_4 = (g_4g_1^{-1})g_1,\\
(R8) \quad & g_3 = (g_3g_2^{-1})g_2,\\
(R9) \quad & g_4 = (g_4g_2^{-1})g_2,\\
(R10) \quad & g_4 = (g_4g_3^{-1})g_3.
\end{aligned}$$

The relations $(R1)$ - $(R4)$ can be evaluated similar to the proof of Lemma 1 and are satisfied due to the equations imposed on the exponents t_{ijk}. For the relations $(R5)$ - $(R10)$ we note that

$$\begin{aligned}
g_4g_1^{-1} &= g_1^{-1}g_4g_5^{-t_{145}},\\
g_3g_1^{-1} &= g_1^{-1}g_3g_4^{-t_{134}}g_5^{t_{134}t_{145}-t_{135}},\\
g_2g_1^{-1} &= g_1^{-1}g_2g_3^{-t_{123}}g_4^{t_{123}t_{134}-t_{124}}g_5^{-t_{123}t_{134}t_{145}+t_{123}t_{135}+t_{124}t_{145}-t_{125}},\\
g_4g_2^{-1} &= g_2^{-1}g_4g_5^{-t_{245}},\\
g_3g_2^{-1} &= g_2^{-1}g_3g_4^{-t_{234}}g_5^{t_{234}t_{245}-t_{235}},\\
g_3g_2^{-1} &= g_2^{-1}g_3g_5^{-t_{235}}.
\end{aligned}$$

These conjugates allow to evaluate the relations $(R5)$-$(R10)$ and thus to determine that the presentation of $G(t)$ is consistent. •

3 Hall polynomials

Next, we consider Hall polynomials for the $\mathcal{T}$-groups of Hirsch length at most 5. We assume that the considered groups are given by consistent presentations of the form $G(t)$ as exhibited in Section 2. We denote $s_2(x) = x(x-1)/2$ and $s_3(x) = x(x-1)(x-2)/6$.

3 Theorem: *Let $t \in \mathbb{Z}^{10}$ so that $G(t)$ is consistent. Then the multiplication in $G(t)$ can be described by Hall polynomials $p_i(a,b)$ with $1 \leq i \leq 5$ and such polynomials are given by*

$$\begin{aligned}
p_1 &= a_1 + b_1,\\
p_2 &= a_2 + b_2,\\
p_3 &= a_3 + b_3 + t_{123}a_2b_1,\\
p_4 &= a_4 + b_4 + t_{124}a_2b_1 + t_{134}a_3b_1 + t_{234}a_3b_2\\
&\quad + t_{123}t_{134}a_2s_2(b_1) + t_{123}t_{234}s_2(a_2)b_1 + t_{123}t_{234}a_2b_1b_2,\\
p_5 &= a_5 + b_5 + t_{345}a_4b_3 + t_{245}a_4b_2 + t_{235}a_3b_2 + t_{145}a_4b_1 + t_{135}a_3b_1 + t_{125}a_2b_1\\
&\quad + t_{234}t_{345}s_2(a_3)b_2 + t_{234}t_{245}a_3s_2(b_2) + t_{134}t_{345}s_2(a_3)b_1 + t_{134}t_{145}a_3s_2(b_1)\\
&\quad + t_{234}t_{345}a_3b_2b_3 + t_{134}t_{345}a_3b_1b_3 + t_{134}t_{245}a_3b_1b_2 + t_{124}t_{345}a_2b_1b_3\\
&\quad + t_{124}t_{345}a_2a_3b_1 + (t_{123}t_{235} + t_{124}t_{245})a_2b_1b_2\\
&\quad + (t_{123}t_{235} + t_{124}t_{245})s_2(a_2)b_1 + (t_{123}t_{135} + t_{124}t_{145})a_2s_2(b_1)\\
&\quad + t_{123}t_{234}t_{245}a_2b_1s_2(b_2) + t_{123}t_{134}t_{245}s_2(a_2)s_2(b_1) + t_{123}t_{234}t_{245}s_3(a_2)b_1\\
&\quad + t_{123}t_{134}t_{145}a_2s_3(b_1) + t_{123}t_{234}t_{245}s_2(a_2)b_1b_2 + t_{123}t_{134}t_{245}a_2s_2(b_1)b_2.
\end{aligned}$$

Proof: We use the approach of Sims [8, page 441ff] to determine the desired polynomials. As a first step, we determine the polynomials $r_{i,j,k}$ with

$$g_i^x g_j^y = g_j^y g_i^x g_{i+1}^{r_{i,j,i+1}(x,y)} \cdots g_n^{r_{i,j,n}(x,y)}$$

for $1 \leq j < i \leq 5$ and all $x, y \in \mathbb{Z}$. Using [7, 5.1.5], we note that the following conditions hold for arbitrary group elements g, h and $x \in \mathbb{Z}$:

$$\begin{aligned}
\text{(1) } [[g,h],g] &= 1 \text{ implies that } [g^x,h] = [g,h]^x,\\
\text{(2) } [[g,h],h] &= 1 \text{ implies that } [g,h^x] = [g,h]^x,\\
\text{(3) } [[g,h],g] &= [[g,h],h] = 1 \text{ implies that } (gh)^x = g^x h^x [g,h]^{s_2(x)}.
\end{aligned}$$

As $[g_4, g_i]$ is central in $G(t)$, we thus obtain that

$$\begin{aligned}
g_4^x g_1^y &= g_1^y g_4^x [g_4^x, g_1^y]\\
&= g_1^y g_4^x [g_4, g_1^y]^x\\
&= g_1^y g_4^x [g_4, g_1]^{xy}\\
&= g_1^y g_4^x g_5^{xyt_{145}},
\end{aligned}$$

$$\begin{aligned}
g_4^x g_2^y &= g_2^y g_4^x g_5^{xyt_{245}},\\
g_4^x g_3^y &= g_3^y g_4^x g_5^{xyt_{345}}.
\end{aligned}$$

Next, we compute the equations for $g_3^x g_1^y$ and $g_3^x g_2^y$ in two steps. First, we show that induction on y yields that

$$(*) \quad g_3^{g_1^y} = g_3 g_4^{yt_{134}} g_5^{yt_{135}+s_2(y)t_{134}t_{145}}.$$

This equation is clearly valid for $y = 0$ and $y = 1$. Assume that it is valid for $y - 1$. Then

$$\begin{aligned}
g_3^{g_1^y} &= (g_3^{g_1^{y-1}})^{g_1} \\
&= (g_3 g_4^{(y-1)t_{134}} g_5^{(y-1)t_{235}+s_2(y-1)t_{134}t_{145}})^{g_1} \\
&= g_3^{g_1} (g_4^{g_1})^{(y-1)t_{134}} (g_5^{g_1})^{(y-1)t_{135}+s_2(y-1)t_{134}t_{145}} \\
&= g_3 g_4^{t_{134}} g_5^{t_{135}} (g_4 g_5^{t_{145}})^{(y-1)t_{134}} g_5^{(y-1)t_{134}+s_2(y-1)t_{134}t_{145}} \\
&= g_3 g_4^{t_{134}} g_5^{t_{135}} g_4^{(y-1)t_{134}} g_5^{(y-1)t_{134}t_{145}} g_5^{(y-1)t_{135}+s_2(y-1)t_{134}t_{145}} \\
&= g_3 g_4^{yt_{134}} g_5^{yt_{135}+s_2(y-1)t_{134}t_{145}+(y-1)t_{134}t_{145}} \\
&= g_3 g_4^{yt_{134}} g_5^{yt_{135}+((y-2)(y-1)/2)t_{134}t_{145}+(y-1)t_{134}t_{145}} \\
&= g_3 g_4^{yt_{134}} g_5^{yt_{135}+(y(y-1)/2)t_{134}t_{135}} \\
&= g_3 g_4^{yt_{134}} g_5^{yt_{135}+s_2(y)t_{134}t_{135}}.
\end{aligned}$$

This proves the desired formula for $y \geq 0$. An analogue computation leads to $g_3^{g_1^y} = g_3 g_4^{yt_{134}} g_5^{yt_{135}+s_2(y)t_{134}t_{145}}$ for $y \leq 0$.
Next note that $(g_3^x)^{g_1^y} = (g_3^{g_1^y})^x$ holds. Hence (3) and $(*)$ yield

$$\begin{aligned}
(g_3^x)^{g_1^y} &= (g_3^{g_1^y})^x \\
&= (g_3 g_4^{yt_{134}} g_5^{yt_{135}+s_2(y)t_{134}t_{145}})^x \\
&= (g_3 g_4^{yt_{134}})^x g_5^{xyt_{135}+xs_2(y)t_{134}t_{145}} \\
&= g_3^x g_4^{xyt_{134}} [g_3, g_4]^{s_2(x)yt_{134}} g_5^{xyt_{135}+xs_2(y)t_{134}t_{145}} \\
&= g_3^x g_4^{xyt_{134}} g_5^{s_2(x)yt_{134}t_{345}+xyt_{135}+xs_2(y)t_{134}t_{145}}.
\end{aligned}$$

Using this and the similar calculation for $g_3^x g_2^y$ we obtain that

$$\begin{aligned}
g_3^x g_1^y &= g_1^y g_3^x g_4^{xyt_{134}} g_5^{s_2(x)yt_{134}t_{345}+xyt_{135}+xs_2(y)t_{134}t_{145}}, \\
g_3^x g_2^y &= g_2^y g_3^x g_4^{xyt_{234}} g_5^{s_2(x)yt_{234}t_{345}+xyt_{235}+xs_2(y)t_{234}t_{245}}.
\end{aligned}$$

We have now determined all polynomials r_{ijk} in the subgroup $G_2 = \langle g_2, \ldots, g_5 \rangle$. Based on this, we can use collection to determine the Hall polynomials of G_2 with this:

$$\begin{aligned}
g^a g^b &= g_2^{a_2} g_3^{a_3} g_4^{a_4} g_5^{a_5} g_2^{b_2} g_3^{b_3} g_4^{b_4} g_5^{b_5} \\
&= g_2^{a_2} g_3^{a_3} g_2^{b_2} g_4^{a_4} g_5^{a_4 b_2 t_{245}} g_5^{a_5} g_3^{b_3} g_4^{b_4} g_5^{b_5} \\
&= g_2^{a_2+b_2} g_3^{a_3} g_4^{a_3 b_2 t_{234}+a_4} g_5^{s_2(a_3)b_2 t_{234}t_{345}+a_3 b_2 t_{235}+a_3 s_2(b_2)t_{134}t_{245}} g_5^{a_4 b_2 t_{245}+a_5} g_3^{b_3} g_4^{b_4} g_5^{b_5} \\
&= g_2^{a_2+b_2} g_3^{a_3+b_3} g_4^{a_3 b_2 t_{234}+a_4} g_5^{b_3 a_3 b_2 t_{234}t_{345}+b_3 a_4 t_{345}} \\
&\quad g_5^{s_2(a_3)b_2 t_{234}t_{345}+a_3 b_2 t_{235}+a_3 s_2(b_2)t_{134}t_{245}+a_4 b_2 t_{245}+a_5} g_4^{b_4} g_5^{b_5} \\
&= g_2^{a_2+b_2} g_3^{a_3+b_3} g_4^{a_3 b_2 t_{234}+a_4+b_4} \\
&\quad g_5^{b_3 a_3 b_2 t_{234}t_{345}+b_3 a_4 t_{345}+s_2(a_3)b_2 t_{234}t_{345}+a_3 b_2 t_{235}+a_3 s_2(b_2)t_{134}t_{245}+a_4 b_2 t_{245}+a_5+b_5}.
\end{aligned}$$

Thus the the multiplication in G_2 is given by

$$g_2^{a_2} \cdots g_5^{a_5} \cdot g_2^{b_2} \cdots g_5^{b_5} = g_2^{f_2(a,b)} \cdots g_5^{f_5(a,b)}$$

with

$$\begin{aligned}
f_2(a,b) &= a_2 + b_2, \\
f_3(a,b) &= a_3 + b_3, \\
f_4(a,b) &= a_4 + b_4 + a_3 b_2 t_{234}, \\
f_5(a,b) &= a_5 + b_5 + a_3 b_2 t_{235} + a_4 b_2 t_{245} + a_4 b_3 t_{345} \\
&\quad + a_3 s_2(b_2) t_{234} t_{245} + s_2(a_3) b_2 t_{234} t_{345} + a_3 b_2 b_3 t_{234} t_{345}.
\end{aligned}$$

Further, these multiplication polynomials allow to determine powering polynomials for G_2. The resulting polynomials with

$$(g_2^{a_2} \cdots g_5^{a_5})^x = g_2^{k_2(a,x)} \cdots g_5^{k_5(a,x)}$$

are given by

$$\begin{aligned}
k_2(a,x) &= f_2(k_2(a,x-1),a) \\
&= k_2(a,x-1) + a_2 \\
&\vdots \\
&= k_2(a,1) + (x-1)a_2 \\
&= x a_2,
\end{aligned}$$

$$\begin{aligned}
k_3(a,x) &= f_3(k_3(a,x-1),a) \\
&= k_3(a,x-1) + a_3 \\
&\vdots \\
&= k_3(a,1) + (x-1)a_3 \\
&= x a_3,
\end{aligned}$$

$$\begin{aligned}
k_4(a,x) &= f_4(k_4(a,x-1),a) \\
&= k_4(a,x-1) + a_4 + k_3(a,x-1) a_2 t_{234} \\
&= k_4(a,x-1) + a_4 + (x-1) a_3 a_2 t_{234} \\
&= k_4(a,x-2) + 2a_4 + (x-1+x-2) a_3 a_2 t_{234} \\
&\vdots \\
&= k_4(a,1) + (x-1)a_4 + ((x-1)x - \sum_{i=1}^{x-1} i) a_3 a_2 t_{234} \\
&= x a_4 + s_2(x) a_2 a_3 t_{234},
\end{aligned}$$

$$k_5(a,x) = f_5(k_5(a,x-1))$$

$$
\begin{aligned}
&= k_5(a,x-1)+a_5+k_3(a,x-1)a_2t_{235}+k_4(a,x-1)a_2t_{245}\\
&\quad +k_4(a,x-1)a_3t_{345}+k_3(a,x-1)s_2(a_2)t_{234}t_{245}\\
&\quad +s_2(k_3(a,x-1))a_2t_{234}t_{345}+k_3(a,x-1)a_2a_3t_{234}t_{345}\\
&= k_5(a,x-1)+a_5+(x-1)a_3a_2t_{235}+((x-1)a_4+s_2(x-1)a_2a_3t_{234})a_2t_{245}\\
&\quad +((x-1)a_4+s_2(x-1)a_2a_3t_{234})a_3t_{345}+(x-1)a_3s_2(a_2)t_{234}t_{245}\\
&\quad +(a_3^2s_2(x-1)+(x-1)s_2(a_3))a_2t_{234}t_{345}+(x-1)a_3a_2a_3t_{234}t_{345}\\
&= k_5(a,x-2)+2a_5+(x-1+x+1)(a_2a_3t_{235}+a_2a_4t_{245}+a_3a_4t_{345})\\
&\quad +(s_2(x-2)+s_2(x-2))(a_2^2a_3t_{234}t_{245}+a_2a_3^2t_{234}t_{345}+a_3^2a_2t_{234}t_{345})\\
&\quad +(x-1+x-2)(s_2(a_3)a_2t_{234}t_{345}+a_2a_3^2t_{234}t_{345}+s_2(a_2)t_{234}t_{345})\\
&= \cdots\\
&\vdots\\
&= k_5(a,1)+(x-1)a_5s_2(x)(a_2a_3t_{235}+a_2a_4t_{245}+a_3a_4t_{345})\\
&\quad +s_3(x)(a_2^2a_3t_{234}t_{245}+a_2a_3^2t_{234}t_{345}+a_3^2a_2t_{234}t_{345})\\
&\quad +s_2(x)(s_2(a_3)a_2t_{234}t_{345}+a_2a_3^2t_{234}t_{345}+s_2(a_2)t_{234}t_{345})\\
&= xa_5+s_2(x)(a_2a_3t_{235}+a_2a_4t_{245}+a_3a_4t_{345})\\
&\quad +s_2(x)a_2a_3t_{234}(t_{245}((2x-1)a_2-3)+t_{345}((4x+1)a_3-3))/6.
\end{aligned}
$$

We now consider the computation for $g_2^x g_1^y$. This is done in two steps. At first we investigate the conjugate $g_2^{g_1^y}$ and write this as $g_2 g_3^{r(y)} g_4^{s(y)} g_5^{t(y)}$ for polynomials $r(y), s(y), t(y)$. We determine these recursively using the polynomials $f_2, \ldots, f_5$:

$$
\begin{aligned}
(g_2)^{g_1^y} &= ((g_2)^{g_1^{y-1}})^{g_1}\\
&= (g_2 g_3^{r(y-1)} g_4^{s(y-1)} g_5^{t(y-1)})^{g_1}\\
&= g_2^{g_1}(g_3^{g_1})^{r(y-1)}(g_4^{g_1})^{s(y-1)}(g_5^{g_1})^{t(y-1)}\\
&= g_2 g_3^{t_{123}} g_4^{t_{124}} g_5^{t_{125}} g_3^{r(y-1)} g_4^{r(y-1)t_{134}} g_5^{s_2(r(y-1))t_{134}t_{345}+r(y-1)t_{135}} g_4^{s(y-1)} g_5^{s(y-1)t_{145}}\\
&\quad g_5^{t(y-1)}\\
&= (g_2 g_3^{t_{123}} g_4^{t_{124}} g_5^{t_{125}})(g_3^{r(y-1)} g_4^{r(y-1)t_{134}+s(y-1)}\\
&\quad g_5^{s_2(r(y-1))t_{134}t_{345}+r(y-1)t_{135}+s(y-1)t_{145}+t(y-1)})\\
&= g_2^{1+0} g_3^{t_{123}+r(y-1)} g_4^{t_{124}+r(y-1)t_{134}+s(y-1)}\\
&\quad g_5^{t_{125}+t(y-1)+s_2(r(y-1))t_{134}t_{345}+r(y-1)t_{135}+s(y-1)t_{145}+t_{124}r(y-1)t_{345}}.
\end{aligned}
$$

We obtain the following conditions:

$$
\begin{aligned}
r(y) &= r(y-1)+t_{123},\\
s(y) &= s(y-1)+t_{124}+r(y-1)t_{134},\\
t(y) &= t(y-1)+t_{125}+s_2(r(y-1))t_{134}t_{345}+r(y-1)(t_{124}t_{345}+t_{135})+s(y-1)t_{145}.
\end{aligned}
$$

Solving these recursions and using that $t_{123}t_{345}=0$ yields

$$
r(y) = r(y-1)+t_{123}
$$

$$
\begin{aligned}
&= r(y-2) + t_{123} + t_{123} \\
&\vdots \\
&= r(1) + (y-1)t_{123} \\
&= yt_{123},
\end{aligned}
$$

$$
\begin{aligned}
s(y) &= s(y-1) + t_{124} + (y-1)t_{123}t_{134} \\
&= s(y-2) + 2t_{124} + (y-1+y-2)t_{123}t_{134} \\
&\vdots \\
&= s(1) + (y-1)t_{124} + ((y-1)y - \sum_{i=1}^{y-1} i)t_{123}t_{134} \\
&= yt_{124} + (y(y-1)/2)t_{123}t_{134} \\
&= yt_{124} + s_2(y)t_{123}t_{134},
\end{aligned}
$$

$$
\begin{aligned}
t(y) &= t(y-1) + t_{125} + (t_{123}^2 s_2(y-1) + (y-1)s_2(t_{123}))t_{134}t_{345} \\
&\quad +(y-1)t_{123}(t_{124}t_{345} + t_{135}) + (y-1)t_{124}t_{145} + s_2(y-1)t_{123}t_{134}t_{145} \\
&= t(y-2) + 2t_{125} + (s_2(y-1) + s_2(y-2))t_{123}^2 t_{134}t_{345} \\
&\quad +(y-1+y-2)s_2(t_{123})t_{134}t_{345} + (y-1+y-2)t_{123}(t_{124}t_{345} + t_{135}) \\
&\quad +(y-1+y-2)t_{124}t_{145} + (s_2(y-1) + s_2(y-2))t_{123}t_{134}t_{145} \\
&= \cdots \\
&\vdots \\
&= t(1) + (y-1)t_{125} \\
&\quad +(y(y-1)/2)(s_2(t_{123})t_{134}t_{345} + t_{123}(t_{124}t_{345} + t_{135}) + t_{124}t_{145}) \\
&\quad +(\sum_{i=1}^{y-1} s_2(i))t_{123}t_{134}(t_{123}t_{345} + t_{145}) \\
&= yt_{125} + s_2(y)(s_2(t_{123})t_{345}t_{134} + t_{123}(t_{124}t_{345} + t_{135}) + t_{124}t_{145}) \\
&\quad +s_3(y)t_{123}t_{134}(t_{123}t_{345} + t_{145}) \\
&= yt_{125} + s_2(y)(t_{123}t_{135} + t_{124}t_{145}) + s_3(y)t_{123}t_{134}t_{145}.
\end{aligned}
$$

Using the polynomials $k_2, \ldots, k_5$ we now determine

$$
g_2^x g_1^y = g_1^y (g_2^{g_1^y})^x = g_1^y (g_2 g_3^{r(y)} g_4^{s(y)} g_5^{t(y)})^x = g_1^y g_2^x g_3^{R(x,y)} g_4^{S(x,y)} g_5^{T(x,y)}
$$

with

$$
\begin{aligned}
R(x,y) &= xr(y) = xyt_{123}, \\
S(x,y) &= xs(y) + s_2(x)r(y)t_{234} \\
&= xyt_{124} + xs_2(y)t_{123}t_{134} + s_2(x)yt_{123}t_{234}, \\
T(x,y) &= xt(y) + s_2(x)(r(y)t_{235} + s(y)t_{245} + r(y)s(y)t_{345}) \\
&\quad +s_2(x)r(y)t_{234}(t_{245}((2x-1)-3) + t_{345}((4x+1)r(y)-3))/6.
\end{aligned}
$$

Thus we have determined all polynomials $r_{i,j,k}$. Based on these, one can use the following GAP program to perform a symbolic collection and thus obtain $p_1, \ldots, p_5$. This program assumes that r and t are global variables so that $r[i][j][k]$ contains the polynomial r_{ijk} and $t[i][j][k]$ is the indeterminate t_{ijk}.

```
CollectSymbolic := function( )
    local a, b, v, stack, c, i, j, k, m;

    # initiate variables
    a := List([1..5],
         x -> Indeterminate(Rationals, Concatenation("a",String(x))));
    b := List([1..5],
         x -> Indeterminate(Rationals, Concatenation("b",String(x))));
    v := StructuralCopy(a);
    v[5] := v[5] + b[5];
    stack := Reversed(List([1..4], x -> [x, b[x]]));

    # perform collection until stack is empty
    while Length(stack) > 0 do
        c := stack[Length(stack)];
        Unbind(stack[Length(stack)]);
        if c[2] <> 0*c[2] then
            i := c[1];
            v[i] := v[i] + c[2];
            for j in Reversed([i+1..4]) do
                for k in Reversed([1..Length(r[i][j])]) do
                    m := StructuralCopy(r[i][j][k]);
                    m[2] := Value(m[2], [x,y], [v[j], c[2]]);
                    if m[1] = 5 then
                        v[5] := v[5] + m[2];
                    elif m[2] <> 0*m[2] then
                        Add(stack, m);
                    fi;
                od;
                v[j] := 0;
            od;
        fi;
    od;

    # divide by consistency relations to simplify polynomial
    v[5] := PolynomialReduction(v[5], [t[1][2][3]*t[3][4][5]],
                                          MonomialLexOrdering())[1];
    return v;
end;
```

•

If G is a $\mathcal{T}$-group of Hirsch length $n < 5$, then Hall polynomials for G can also be read off from Theorem 3: These are given by $p_1, \ldots, p_n$.

References

[1] B. Eick. Algorithms for polycyclic groups. Habilitationsschrift, Universität Kassel, 2001.

[2] B. Eick and A.-K. Engel. The torsion free nilpotent groups of hirsch length at most 5. *Submitted*, 2016.

[3] P. Hall. *The Edmonton notes on nilpotent groups.* Queen Mary College Mathematics Notes. Mathematics Department, Queen Mary College, London, 1969.

[4] P. Hall. *The collected works of Philip Hall.* Oxford Science Publications. The Clarendon Press Oxford University Press, New York, 1988. Compiled and with a preface by K. W. Gruenberg and J. E. Roseblade, With an obituary by Roseblade.

[5] D. F. Holt, B. Eick, and E. A. O'Brien. *Handbook of computational group theory.* Discrete Mathematics and its Applications (Boca Raton). Chapman & Hall/CRC, Boca Raton, FL, 2005.

[6] C. R. Leedham-Green and L. H. Soicher. Symbolic collection using Deep Thought. *LMS J. Comput. Math.*, 1:9 – 24, 1998.

[7] D. J. S. Robinson. *A course in the theory of groups.* Springer, Graduate texts in mathematics 80, 1982.

[8] C. C. Sims. *Computation with finitely presented groups.* Cambridge University Press, Cambridge, 1994.